

Single production of charged gauge bosons from little Higgs models in association with top quark at the *LHC*

Chong-Xing Yue, Shuo Yang, Li-Hong Wang

Department of Physics, Liaoning Normal University, Dalian 116029, China^{*}

February 2, 2008

Abstract

In the context of the little Higgs models, we discuss single production of the new charged gauge bosons in association with top quark at the *CERN* Large Hadron Collider(*LHC*). We find that the new charged gauge bosons W_H^- and X^- , which are predicted by the littlest Higgs model and the $SU(3)$ simple model, respectively, can be abundantly produced at the *LHC*. However, since the main backgrounds coming from the processes $pp \rightarrow t\bar{t} + X$ and $pp \rightarrow tW^- + X$ are very large, the values of the ratios N_W and N_X are very small in most of the parameter space. It is only possible to detect the signal of the gauge boson W_H^- via the process $pp \rightarrow gb + X \rightarrow tW_H^- + X$ at the *LHC* in a small region of the parameter space.

PACS number(s): 12.60.Cn, 14.65.Ha, 12.15.-y

^{*}E-mail: cxyue@lnnu.edu.cn

I. Introduction

Most of the new physics models predict the existence of the new gauge bosons, generally called Z' or W' with masses in the TeV range. If these new particles are discovered, they would represent irrefutable proof of new physics, most likely that the gauge group of the standard model (SM) must be extended. Thus, search for extra gauge bosons provides a common tool in quest for new physics at the next generation collider experiments[1]. The discovery and study of extra gauge bosons is one of the important goals of the *CERN* Large Hadron Collider (*LHC*).

The little Higgs models[2] propose a new approach to electroweak symmetry breaking ($EWSB$) accomplished by a naturally light Higgs sector. These models predict new gauge bosons, fermions, and scalars at or below the TeV scale, which might generate characteristic signatures at the next generation collider experiments, especially at the *LHC*[3,4]. The little Higgs models generally predict the existence of the pure left-handed charged gauge boson W' , which has the SM -like couplings to ordinary particles and might generate significant corrections to single top quark production at the *LHC*[5]. It has been shown that the charged gauge bosons from little Higgs models should be either seen or excluded in the first year of running at the *LHC*[6]. Thus, it is very interesting to study production of these new charged gauge bosons at the *LHC*.

In the SM framework, single production of the charged gauge bosons $W^\pm$ associated with a top quark is one of the important process for single top quark production. Its cross section is negligible at the *Tevatron* but of considerable size at the *LHC*, where it is larger than that of the s -channel process for single top quark production. There is an extensive work on tW associated production at the *LHC* in the literature[7,8,9]. The production cross section for this process has been calculated at leading order, with a subset of the next-to-leading order (NLO) corrections included[7]. Including both the subsequent leptonic decays $W \rightarrow l\nu$ and $t \rightarrow l\nu b$, as well as the emission of real radiation in the top decay, a NLO calculation of this process has been given in Ref.[8]. Recently, Ref.[9] gives a complete calculation of electroweak supersymmetric effects at one loop to tW associated production at the *LHC*. In this letter, we will study single production

of the charged gauge bosons predicted by the little Higgs models in association with top quark and see whether the possible signals of these new particles can be detected via this process at the *LHC*.

II. The relevant formulas

Based on the structure of the extended electroweak gauge group, the little Higgs models can be generally divided into two classes[3,10]: the product group models in which the $SU(2)_L$ gauge group arises from the diagonal breaking of two or more gauge groups, and the simple gauge group models in which the $SU(2)_L$ gauge group arises from the breaking of a single larger gauge group down to an $SU(2)$ subgroup. The littlest Higgs models (*LH*)[11] and the $SU(3)$ simple group model[10,12] are the simplest examples of the product group models and the simple group models, respectively. All of these little Higgs models predict the pure left-handed charged gauge boson W' . The couplings of the new particle W' to fermions can be unitive written as:

$$\mathcal{L}_{W'q_iq_j} = \frac{eA}{\sqrt{2}S_W} V_{ij} W'_\mu \bar{q}_i \gamma^\mu P_L q_j, \quad (1)$$

where $P_L = (1 - \gamma_5)/2$, V_{ij} is the *CKM* matrix element, and $S_W = \sin \theta_W$ (θ_W is the Weinberg angle). For the charged gauge boson W_H^- predicted by the *LH* model, the constant A equals to c/s for $(q_i, q_j) = (u, d), (c, s)$, and (t, b) , in which $c(s = \sqrt{1 - c^2})$ is the mixing parameter between $SU(2)_1$ and $SU(2)_2$ gauge bosons[13]. For the charged gauge boson X^- predicted by the $SU(3)$ simple group model, the constant A equals to δ_t and δ_ν for $(q_i, q_j) = (t, b)$ and (u, d) [or (c, s)], respectively. The factors δ_t and δ_ν can be written as[3]:

$$\delta_t = \frac{\nu}{\sqrt{2}f} t_\beta \frac{x_\lambda^2 - 1}{x_\lambda^2 + t_\beta^2}, \quad \delta_\nu = -\frac{\nu}{2ft_\beta}. \quad (2)$$

Where $\nu = 246 \text{ GeV}$, $f = \sqrt{f_1^2 + f_2^2}$, $t_\beta = \tan \beta = f_2/f_1$, and $x_\lambda = \lambda_1/\lambda_2$, in which f_1 and f_2 are the vacuum condensate values of the two sigma-model fields Φ_1 and Φ_2 , respectively. λ_1 and λ_2 are the *Yukawa* coupling parameters.

It has been shown that the collinear $\bar{b}$ component of the inclusive process $gg \rightarrow t\bar{b}W^-$ can be attached as a *QCD NLO* correction to bottom quark distribution function of the

exclusive process $bg \rightarrow tW^-$ at the partonic level[7,8,9]. The LO contributions to tW^- associated production is best considered to arise from the process $bg \rightarrow tW^-$. Thus, in this paper, we only consider single production of the heavy charged gauge boson W_i^- in association with a top quark via the two body final state process $bg \rightarrow tW_i^-$ at the LHC as shown in Fig.1 in which $i = 1$ and 2 represent the gauge bosons W_H^- and X^- , respectively. Our numerical results are easy transferred to those for single production of the heavy gauge boson W_i^+ at the LHC by replacing $\bar{b}$ as b and $\bar{t}$ as t .




Figure 1: Feynman diagrams for the partonic process $bg \rightarrow tW_i^-$

At the leading order, the scattering amplitude of the partonic process $bg \rightarrow tW_i^-$ can be written as:

$$M = \frac{eg_s A}{\sqrt{2}S_W} \bar{u}(t) \left[\frac{\not{\epsilon}_2 P_L (\not{P}_g + \not{P}_b + m_b) \not{\epsilon}_1}{\hat{s}' - m_b^2} + \frac{\not{\epsilon}_1 (\not{P}_t - \not{P}_g + m_t) \not{\epsilon}_2 P_L}{\hat{u} - m_t^2} \right] u(b), \quad (3)$$

where $\hat{s}' = (P_g + P_b)^2 = (P_{W_i} + P_t)^2$, $\hat{u} = (P_t - P_g)^2 = (P_b - P_{W_i})^2$. ϵ_1 and ϵ_2 are the gluon and W_i^- polarization vectors, respectively.

The cross section for single production of the heavy gauge boson $W_i^- = W_H^-$ or X^- associated with a top quark at the LHC with the center-of-mass $\sqrt{S} = 14TeV$ can be obtained by convoluting the production cross section $\hat{\sigma}(tW_i^-)$ of the partonic process $gb \rightarrow tW_i^-$ with the parton distribution functions ($PDF's$):

$$\sigma_i(S) = \int_{\tau}^1 dx_1 \int_{\tau/x_1}^1 dx_2 [f_{g/p}(x_1, \mu) f_{b/p}(x_2, \mu) \hat{\sigma}(tW_i^-) + f_{b/p}(x_1, \mu) f_{g/p}(x_2, \mu) \hat{\sigma}(tW_i^-)], \quad (4)$$

where $\tau = (M_{W_i}^2 + m_t^2)/S$ and $\hat{s}' = x_1 x_2 S$. Through out this paper, we will use $CTEQ6L$ $PDF's$ [14] for the bottom quark and gluon $PDF's$. Following the suggestions given by

Refs.[7,8,9], we assume that the factorization scale μ for the bottom quark *PDF* is of order $(M_{W_i} + m_t)/4$.

III. Numerical results and conclusions

Except for the *SM* input parameter $m_t = 171.4\text{GeV}$ [15], $\alpha_e = 1/128.8$, $\alpha_s = 0.118$, and $S_W^2 = 0.2315$ [16], the production cross sections of the new charged gauge bosons W_H^- and X^- coming from the *LH* model and the *SU*(3) simple group model are dependent on the free parameters (M_{W_H}, c) and $(M_X, x_\lambda, t_\beta)$, respectively. Considering the constraints of the electroweak precision data on these free parameters, we will assume $1\text{TeV} \leq M_{W_H} \leq 2\text{TeV}$ and $0 < c \leq 0.6$ for the *LH* model[17] and $1\text{TeV} \leq M_X \leq 2\text{TeV}$, $x_\lambda > 1$, and $t_\beta > 1$ for the *SU*(3) simple group model[3,10,12] in our numerical calculation.




Figure 2: The cross section σ_W as a function of the mass parameter M_{W_H} for the mixing parameter $c=0.2$ (solid line), 0.4 (dashed line), and 0.6 (dashed-dotted line).

The cross sections σ_W and σ_X for single production of the new gauge bosons W_H^- and X^- associated with top quark at the *LHC* with $\sqrt{S} = 14\text{TeV}$ are plotted as functions of the mass parameters M_{W_H} and M_X in Fig.2 and Fig.3, respectively, in which we have taken different values of the free parameters c , x_λ , and t_β . One can see from these figures that the single production cross section for the gauge boson W_H^- is larger than that for the gauge boson X^- , which is because compared to the coupling $W_H^- t \bar{b}$, the coupling $X^- t \bar{b}$ is suppressed by the factor ν/f . For $0.2 \leq c \leq 0.6$, $1\text{TeV} \leq M_{W_H} \leq 2\text{TeV}$, the value of

the production cross section σ_W is in the range of $0.7fb \sim 286.7fb$. The production cross section σ_X increase as the free parameter x_λ increasing. For $x_\lambda = 5$, $1 \leq t_\beta \leq 3$, and $1TeV \leq M_X \leq 2TeV$, the value of σ_X is in the range of $0.16fb \sim 15.3fb$.




Figure 3: The cross section σ_X as a function of the mass parameter M_X for different values of the free parameters x_λ and t_β .

To see whether the heavy gauge bosons W_H^- and X^- can be observed at the *LHC* via the processes $pp \rightarrow gb + X \rightarrow tW_H^- + X$ and $pp \rightarrow gb + X \rightarrow tX^- + X$, we discuss the possible decay modes of the heavy gauge bosons W_H^- and X^- . For W_H^- , the main decay modes are W^-H , $\bar{q}q'$, and $l^-\nu$, in which $q(q')$ and l^- present all three generation quarks and leptons, respectively. If we neglect the final state masses, then we have $\Gamma(W_H^- \rightarrow \bar{q}q') = 3\Gamma(l^-\nu)$. At leading order, the total decay width $\Gamma_{W_H^-}$ can be approximately written

as[17]:

$$\Gamma_{W_H^-} = \frac{\alpha_e}{96S_W^2} \left[\frac{96c^2}{s^2} + \frac{C_W^2(c^2 - s^2)^2}{s^2c^2} \right] M_{W_H}. \quad (5)$$

For the heavy gauge boson X^- , the main decay modes are $\bar{q}q'$ and $l^-\nu$. The total decay width Γ_{X^-} can be approximately written as[3]:

$$\Gamma_{X^-} = \frac{\alpha_e M_X}{4S_W^2} (\delta_t^2 + 5\delta_\nu^2). \quad (6)$$

In general, to reject backgrounds in hadronic collider environment, the heavy gauge bosons may be most likely to be observed via their pure leptonic decays. In this case, single production of the heavy gauge boson W_H^- or X^- associated with a top quark gives the possible observable five fermion final states with at least one b quark $\nu\nu l^+ l^- b$ with $W_H^-(X^-) \rightarrow l^-\nu$ and $t \rightarrow W^+ b \rightarrow \nu l^+ b$. The backgrounds of this kind signals mainly come from the SM processes $pp \rightarrow t\bar{t} + X$ and $pp \rightarrow tW^- + X$, in which for the $t\bar{t}$ production process, one of the bottom quarks from a top decay is assumed missing detection[7]. Under narrow width approximation, the number of this kind signal events can be written as $S_W = \mathcal{L}_{int} \sigma_W Br(W_H^- \rightarrow l^-\nu) Br(t \rightarrow \nu l^+ b)$ and $S_X = \mathcal{L}_{int} \sigma_X Br(X^- \rightarrow l^-\nu) Br(t \rightarrow \nu l^+ b)$, in which $\mathcal{L}_{int}$ is the yearly integrated luminosity of the LHC with $\sqrt{s} = 14 TeV$. In our numerical estimation, we will take $\mathcal{L}_{int} = 100 fb^{-1}$.




Figure 4: The ratio of signal over square root of the background ($S_W/\sqrt{B}$) for the W_H^- production associated with top quark at the LHC .

To see the signals against the backgrounds for the heavy gauge bosons W_H^- and X^- , we introduce the ratio of signal over square root of the backgrounds: $N_W = S_W/\sqrt{B}$ and $N_X = S_X/\sqrt{B}$, in which B includes the contributions of $t\bar{t}$ production process and tW^- production process. To simple our calculation, we assume that all charged gauge bosons W_H^- , X^- and W^+ decay to $e\nu_e$. Similar to Ref.[8], we take the appropriate cuts on the transverse momentum $P_T(e)$ and rapidly $\eta(e)$ for the final electron as: $P_T(e) > 20GeV$ and $|\eta(e)| \leq 2.5$. For the b quark, which comes from top quark decay, we take the cuts as: $P_T(b) > 30GeV$ and $|\eta(b)| \leq 3$. Our numerical result are shown in Fig.4, in which we plot the ratio N_W as a function of the mass M_{W_H} for three values of the free parameter c . Our numerical results show that the value of the ratio N_X is smaller than 0.1 in most of the parameter space of the $SU(3)$ simple model. So we do not give our numerical results for the heavy gauge boson X^- in Fig.4. One can see from Fig.4 that for $c \geq 0.5$ and $M_{W_H} \leq 1.5TeV$, the value of the ratio N_W is larger than 1, which might be detected at the LHC [1].

Little Higgs models can be generally divided in two classes: product group models and simple group models. All of the little Higgs models predict the existence of the new charged gauge boson W' with mass in TeV range, which has the SM -like couplings to the ordinary fermions. In this letter, we investigate single production of these new gauge bosons associated with a top quark at the LHC . We find that the new charged gauge bosons W_H^- and X^- , which are predicted by the LH model and the $SU(3)$ simple model, respectively, can be abundantly produced at the LHC . However, since the main backgrounds coming from the processes $pp \rightarrow t\bar{t} + X$ and $pp \rightarrow tW^- + X$ are very large, the values of the ratios N_W and N_X are very small in most of the parameter space. The possible signals of the gauge bosons X^- can not be observed via the process $pp \rightarrow gb + X \rightarrow tX^- + X$ at the LHC . For the gauge boson W_H^- , it might be possible to detect its signal via the process $pp \rightarrow gb + X \rightarrow tW_H^- + X$ at the LHC only for the mixing parameter $c \geq 0.5$ and the mass parameter $M_{W_H} \leq 1.5TeV$.

Acknowledgments

C. X. Yue would like to thank the **Abdus Salam** International Centre for Theoretical Physics(ICTP) for partial support. This work was supported in part by Program for New Century Excellent Talents in University(NCET-04-0290), the National Natural Science Foundation of China under the Grants No.10475037.

References

- [1] G. Weiglein et al.[*LHC/LC* study Group], *hep-ph/0410364*.
- [2] For recent review see: M. Schmaltz and D. Tucker-Smith, *Ann. Rev. Nucl. Part. Sci.* **55**(2005)299.
- [3] T. Han, H. E. Logan, and L. T. Wang, *JHEP* **0601**(2006)099.
- [4] M. Perelstein, *hep-ph/0512128*.
- [5] Chong-Xing Yue, Li Zhou, Shuo Yang, *hep-ph/0604001*.
- [6] Z. Sullivan, *hep-ph/0306266*.
- [7] A. P. Heinson, A. S. Belyaev, E. E. Boos, *Phys. Rev. D***56**(1997)3114; T. M. P. Tait, *Phys. Rev. D***61**(2000)034001; A. Belyaev, E. Boos, *Phys. Rev. D***63**(2001)034012.
- [8] J. Campbell and F. Tramontano, *Nucl. Phys. B***726**(2005)109.
- [9] M. Beccaria, F. M. Denard, C. Verzegnassi, *Phys. Rev. D***71**(2005)033005; M. Beccaria, G. Macorini, F. M. Renard and C. Verzegnassi, *Phys. Rev. D***73**(2006)093001.
- [10] D. E. Kaplan and M. Schmaltz, *JHEP* **0310**(2003)039.
- [11] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, *JHEP* **0207**(2002)034.
- [12] M. Schmaltz, *JHEP* **0408**(2004)056.

- [13] T. Han, H. E. Logan, B. McElrath and L. T. Wang, *Phys. Rev. D***67**(2003)095004.
- [14] J. Pumplin *et al.*, *JHEP* **0207**(2002)012; D. Stump *et al.*, *JHEP* **0310**(2003)046.
- [15] The Tevatron Electroweak Working Group for CDF and D0 Collaborations, *Combination of CDF and D0 Results on the Mass of the Top Quark* , *hep-ex/0608032*
- [16] S. Eiddman *et al.*[Particle Data Gruop], *Phys. Lett. B***592**(2004)1.
- [17] R. Casalbuoni, A. Deandrea, M. Oertel, **JHEP** **0402**(2004)032; Mu-Chun Chen and S. Dawson, *Phys. Rev. D***70**(2004)015003; Chong-Xing Yue and Wei Wang, *Nucl. Phys. B***683**(2004)48; W. Kilian and J. Reuter, *Phys. Rev. D***70**(2004)015004; A. Deandrea, *hep-ph/0405120*; J. L. Hewett, F. J. Petriello and T. G. Rizzo, *JHEP* **0310**(2003)062; C. Csaki, J. Hubisz, G. D. Kribs, P. Meade, and J. Terning, *Phys. Rev. D***67**(2003)115002; C. Csaki et al., *Phys. Rev. D***68**(2003)035009; T. Gregoire, D. R. Smith and J. G. Wacker, *Phys. Rev. D***69**(2004)115008; Mu-chun Chen, *Mod. Phys. Lett. A***21**(2006)621.